

One-shot decoupling and Page curves from a dynamical model for black hole evaporation

Kamil Brádler*

*Department of Mathematics and Statistics, University of Ottawa, Ottawa, Canada
Max Planck Centre for Extreme and Quantum Photonics, University of Ottawa, Ottawa, Canada and
Department of Astronomy and Physics, Saint Mary's University, Halifax, Canada*

Christoph Adami†

Department of Physics and Astronomy, Michigan State University, East Lansing, MI 48824

One-shot decoupling is a powerful primitive in quantum information theory and was hypothesized to play a role in the black hole information paradox. We study black hole dynamics modeled by a trilinear Hamiltonian whose semiclassical limit gives rise to Hawking radiation. An explicit numerical calculation of the discretized path integral of the S -matrix shows that decoupling is exact in the continuous limit, implying that quantum information is perfectly transferred from the black hole to radiation. A striking consequence of decoupling is the emergence of an output radiation entropy profile that follows Page's prediction. We argue that information transfer and the emergence of Page curves is a robust feature of any multi-linear interaction Hamiltonian with a bounded spectrum.

The explicit mechanism of black hole unitary evaporation is a major open problem in high-energy physics [1]. The evaporation of a macroscopic black hole is believed to be well described by a semiclassical result of Hawking [2], but it has also been argued that for a sufficiently old black hole the in-falling Hawking quanta (hereafter called the “partner modes”, as opposed to the radiated modes that constitute the Hawking radiation) must somehow resurface in order to preserve the manifest unitarity of the black hole evolution. Page [3] argued for the latter using a description based entirely on calculating the entropy of the Hawking radiation as a subsystem of the joint pure state of a black hole and its radiation field, without proposing an explicit interaction that would enable this. In the modern language of quantum information theory, Page's calculation implies that when the black hole entropy vanishes, it is *decoupled* from its surroundings. Here we present a model in which black hole modes interact with radiation modes both inside and outside of the horizon in a fully unitary manner, and that allows us to explicitly calculate the entropy of the black hole over time as it gradually decouples from its surroundings.

The modern approach to the resolution of the problem started with Ref. [4]. These authors proposed an abstract mechanism for black hole evaporation based on the recently developed “decoupling” framework (see a general version in [5]). Decoupling theory is a fundamental technique to prove a plethora of quantum communication protocols such as state-merging [6], the fully quantum Slepian-Wolf (FQSW) protocol [7] or the channel capacity coding theorem [8].

The central result of the one-shot decoupling theorem [9] implies the existence of a completely positive (CP) map $\hat{\mathcal{T}}_{A \rightarrow B}$ and a state ρ_{RA} where the following

condition is satisfied

$$\|\hat{\mathcal{T}}_{A \rightarrow B}(U\rho_{RA}U^\dagger) - \vartheta_B \otimes \rho_R\|_1 \rightarrow 0 \quad (1)$$

for almost all unitaries U . Here, $\|A\|_1 \stackrel{\text{def}}{=} \text{Tr}[(A^\dagger A)^{1/2}]$ is the trace norm, R is an auxiliary system, $\rho_R = \text{Tr}_A[\rho_{RA}]$ and ϑ_B is a state specified later. For our purposes it will turn out to be sufficient to consider ρ_{RA} as a maximally entangled state (denoted Φ_{RA}). Then, decoupling Eq. (1) is equivalent to the conditions of the entanglement of the A subsystem of Φ_{RA} being transferred to E via another CP map $\mathcal{T}_{A \rightarrow E}$ called a *complementary* channel to $\hat{\mathcal{T}}_{A \rightarrow B}$.

In what follows we examine decoupling induced by a boson trilinear operator describing the interaction of a quantized black hole's modes with those of radiation inside and outside of the event horizon, in the special case when $\mathcal{T}_{A \rightarrow E}$ is an identity, something that, crucially, is not an assumption of the model but rather an unexpected and natural consequence of the interaction Hamiltonian we use.

In the semiclassical description, the outgoing Hawking radiation shrinks the black hole but the exact mechanism of how the black hole loses its mass over time is beyond the scope of Hawking's original derivation, as in that approach states from past infinity are mapped to future infinity via a Bogoliubov transformation. In order to restore an explicit time-dependence to the evaporation process, we can use the by now well-known insight [10, 11] (see also [12, 13]) that any solution to a time-dependent quantum harmonic oscillator can be understood in terms of a Bogoliubov transformation that connects the initial to the final frequency, and in particular can be implemented via a time-dependent Hamiltonian of the “squeezing” type in which the coupling strength $r(t)$ plays the role of the classical time-dependent driving field (we set

$\hbar = 1$)

$$H = ir(t)(b^\dagger c^\dagger - bc) . \quad (2)$$

Here, the annihilation operators b and c annihilate radiation modes within and outside of the event (or apparent) horizon, respectively. Such a description can be used to derive Hawking radiation from a collapsing shell of matter, for example [14–16]. However, in such a time-dependent semi-classical description, the black hole degrees of freedom are not explicitly quantized and the backreaction of the radiation on the black hole cannot be followed.

The dynamic model for the evaporation of a black hole that we analyze here can be seen as an extension of Hawking’s semi-classical approach [17] where the black hole internal degrees of freedom are explicitly quantized to allow us to track the black hole quantum entropy over time. This model is explicitly *ad hoc*: the quantum interaction is not derived from a quantum field theory that couples gravitational modes to radiation modes. Rather, it is an extension of the semi-classical Hamiltonian (2), and reduces to it when the quantum black hole modes are replaced by their classical expectation values. In the static infinite-time limit, the quantum Hamiltonian (shown below) therefore implements the standard Bogoliubov transformation between past and future infinity, and reproduces the celebrated Hawking results as has been shown previously [11, 18, 19].

The Hamiltonian we study is the so-called boson trilinear Hamiltonian

$$H_{\text{tri}} = ir(ab^\dagger c^\dagger - a^\dagger bc) \quad (3)$$

often encountered in quantum optics and condensed matter physics [20–22] or quantum optomechanics [23]. In quantum optics, the operators b and c refer to the signal and idler modes, respectively, while a refers to the pump mode. The intriguing formal analogy between black hole radiance and parametric amplification has been noted before [18, 19, 24]. As a model of black hole and radiation interactions, the operator a in (3) annihilates black hole modes, whereas c and b are defined as in (2).

The black hole mass is a function of the input state $n_a = \langle a^\dagger a \rangle$ as well as of the coupling strength r (called the *gain* or *squeezing* parameter in quantum optics). In the semi-classical limit, the relationship between $r(t)$ in (2) and black hole parameters is simply (in units where $\hbar = c = G = k = 1$) [18]

$$T(t) = \frac{\omega}{2 \ln[\tanh^{-1}(r(t)t)]} \quad (4)$$

where $T(t)$ is the time-dependent Hawking temperature $T(t) = (8\pi M(t))^{-1}$ and $M(t)$ is the black hole mass.

The initial black hole state can be pure or mixed. For simplicity, we restrict our analysis here to the pure state scenario and a black hole in initial state $|n\rangle_a$ (n modes),

but our results naturally carry over to an arbitrary superposition and any mixed initial black hole state with corresponding nonzero initial entropy as we discuss below. We now define the time-evolution operator on a quantum state $|\Psi(t_1)\rangle$ as $|\Psi(t_2)\rangle = U(t_2, t_1)|\Psi(t_1)\rangle$ where

$$U(t_2, t_1) = \mathcal{T} e^{-i \int_{t_1}^{t_2} H(t') dt'} . \quad (5)$$

The symbol \mathcal{T} stands for Dyson’s time-ordering operator and $H(t)$ is the Hamiltonian describing the unitary evolution of the quantum state. We can then write the time-evolution of the black hole initial state $|\Psi(0)\rangle = |n\rangle_a |0\rangle_b |0\rangle_c \equiv |n\rangle_a |0\rangle_{bc}$ as

$$|\Psi(t)\rangle = U(t, 0) |\Psi(0)\rangle = \mathcal{T} e^{-i \int_0^t H_{\text{tri}}(t') dt'} |n\rangle_a |0\rangle_{bc}, \quad (6)$$

where H_{tri} is the trilinear Hamiltonian (written as a sum over modes)

$$H_{\text{tri}} = \sum_{k=-\infty}^{\infty} ir_{\omega_k}(t)(a_k b_k^\dagger c_k^\dagger - a_k^\dagger b_k c_k). \quad (7)$$

In the following, we will focus on a single mode k with energy ω_k , and omit the index k for convenience.

To calculate (6), we introduce small time slices Δt and discretize the path integral so that with $t = N\Delta t$

$$U(t) = \mathcal{T} e^{-i \int_0^t H_{\text{tri}}(t') dt'} \approx \prod_{i=1}^N e^{-i \Delta t H_i}, \quad (8)$$

where the i -th time-slice Hamiltonian $H_i = ir_0(ab_i^\dagger c_i^\dagger - a_i^\dagger b_i c_i)$ acts on the black hole state and the i -th slice of the bc Hilbert space $|0\rangle_{b_i c_i}$. The initial value r_0 sets the energy scale, and we simply set $r_0 = 1$ in the following.

We now apply (8), so that

$$|\Psi(t)\rangle_{a'bc} = \mathcal{W} |n\rangle_a |0\rangle_{bc} = \prod_{i=1}^N e^{-i \Delta t H_i} |n\rangle_a |0\rangle_{bc}, \quad (9)$$

where $|0\rangle_{bc} \stackrel{\text{df}}{=} |0\rangle_{b_N c_N} \otimes \dots \otimes |0\rangle_{b_1 c_1}$, $W^{(i)} = e^{-i \Delta t H_i}$ and $\mathcal{W} = W^{(N)} \circ \dots \circ W^{(1)}$. The first time slice is

$$|\Psi(1)\rangle = W^{(1)} |n\rangle_a |0\rangle_{b_1 c_1} = \sum_{j=0}^n U_{nj}^{(1)} |j\rangle_a |n-j\rangle_{b_1 c_1} \quad (10)$$

while $|\Psi(t)\rangle$ is given by

$$|\Psi(t)\rangle_{a'bc} = \sum_{j_1 \dots j_N} U_{nj}^{(1)} \dots U_{j_{N-1} j_N}^{(N)} \times |j_N\rangle_a |j_{N-1} - j_N\rangle_{b_N c_N} \dots |n - j_1\rangle_{b_1 c_1}. \quad (11)$$

Note that approximating the path integral using a single time-slice—the static path approximation (SPA), see e.g. [25, 26]—can yield good results at very low temperatures, but disregards self-consistent temporal fluctuations. Indeed, SPA calculations of the black hole entropy using the trilinear Hamiltonian lead to an oscillating behavior of the black hole entropy [18, 19]. The black




FIG. 1. The action of the discretized path integral $W^{(i)}$ is sketched. The braces indicate the entropy calculation of the radiated modes at each time-slice.

hole von Neumann entropy can be calculated using the marginal density matrix $\vartheta_{a'}(t) = \text{Tr}_{bc} [|\Psi(t)\rangle\langle\Psi(t)|_{a'bc}]$, which can be written entirely in terms of the coefficients $|U_{ij}|^2$, which describe the probability to observe j modes of the black hole given that in the previous iteration we found i modes) $\vartheta_{a'}(t) = \sum_{j_N=0}^n p_{j_N} |j_N\rangle\langle j_N|$ where

$$p_{j_N} = \sum_{j_1 > j_2 > \dots > j_{N-1}}^n |U_{n j_1}^{(1)}|^2 \dots |U_{j_{N-2} j_{N-1}}^{(N-1)}|^2 |U_{j_{N-1} j_N}^{(N)}|^2. \quad (12)$$

To follow the time evolution of the black hole entropy $H_n(t) = -\text{Tr} [\vartheta_{a'}(t) \log \vartheta_{a'}(t)]$ given the initial state of the black hole is a pure state with n quanta, we need to evaluate the n coefficients p_{j_N} above. This is a difficult calculation because unlike the Hawking isometry, the unitary operator $W^{(i)}$ is not known to be factorizable in a simple way. The usual formal factorization formulas [27–29] are infinite operator products and often are not suitable for practical calculations. A common route to obtain the time evolution is to expand the exponential around $r_0 t \approx 0$ but this allows only a short-time analysis of the evolution operator. For greater $r_0 t$, the Taylor expansion is prohibitively inefficient. However, a method developed in Ref. [?] enables us to find the action of the unitary operators even for large values of $r_0 t$. The action of $W^{(i)}$ acquires an interesting combinatorial interpretation in terms of an integer lattice known as a generalized Dyck path [30] if it acts on any state generated by the repeated action of $ab^\dagger c^\dagger$ on a ground state $|\text{gr}\rangle$, defined by $a^\dagger bc |\text{gr}\rangle = 0$. The basis elements $\{|k 00\rangle_{abc}\}_{k=0}^n$ spanning our input Hilbert space of $W^{(i)}$ are all ground states of H_i .

We evaluated the output entropy for black holes with initial quanta n for $0 \leq n \leq 50$ using the discretized path integral (11). The resulting entropy as a function of the number of time slices (at constant small Δt) is shown in Fig. 2 for $n = 5, 20$ and 50 . An entropy curve strikingly similar to the one predicted by Page [3] emerges in the limit of large t (several thousand time slices). This shape is consequence of the fact that the final black hole den-

sity matrix $\vartheta_{a'}$ (for $N \gg 0$) is extremely close to a pure vacuum state $|0\rangle_{a'}$ for all examined input basis states $|n\rangle$. Note that the case $n = 50$ gives rise to an extremely large Hilbert space, and to obtain a reliable Taylor expansion for $\Delta t = 1/15$ would require expansion to order up to 500. A straightforward numerical evaluation of the 2^{500} terms is of course impossible, but the method developed in [?] makes the calculation tractable on a High-Performance Computing Cluster. We have also performed a calculation of the integral where partner modes are allowed to interact with the black hole a second time, breaking the symmetry between partner and Hawking modes. Such a calculation is far more complex, but does not change the shape of the Page curves appreciably.

It is important to stress that even though we studied the effect of \mathcal{W} only on basis states $|n\rangle$, we can extend this conclusion to an arbitrary input pure state $\psi_a = \sum_{k=0}^n \beta_k |k\rangle_a$. It is sufficient to show decoupling of the spanning basis set. Thus, ultimately the action of \mathcal{W} can be reformulated as inducing

$$|k\rangle_a |0\rangle_{bc} \xrightarrow{\mathcal{W}} |\Psi(t)\rangle_{a'bc} \approx |0\rangle_{a'} \otimes \vartheta_{bc}(|k\rangle), \quad (13)$$

where $0 \leq k \leq n$ and $\vartheta_{bc}(|k\rangle) \stackrel{\text{df}}{=} \text{Tr}_{a'} [|\Psi(t)\rangle\langle\Psi(t)|_{a'bc}]$. Hence $\vartheta_{bc}(|k\rangle)$ is necessarily pure and mutually orthogonal for all k . Since $|0\rangle_a$ is a “constant” state (independent of k), it immediately follows that an arbitrary superposition ψ_a will be transferred to the Hawking modes and their partners (bc). These are precisely the circumstances captured by the decoupling theorem in Eq. (1) upon identifying $A \Leftarrow a, E \Leftarrow bc$ and $B \Leftarrow a'$. The presence of an input bipartite state Φ_{RA} instead of ψ_A is just a convenient reformulation so that if the decoupling (transfer) of Φ_{RA} is successful, it can be used to perfectly teleport any ψ_A .

As is implied by (13), the black hole evolution ultimately turns out to be an *erasure map* $\hat{\mathcal{T}}_{a \rightarrow a'} : \psi_a \rightarrow \text{Tr}[\psi_a] |0\rangle_{a'}$, which we interpret as complete evaporation. This immediately implies that the map $\mathcal{T}_{a \rightarrow bc}$ must be the identity as announced earlier. The decoupling theorem then implies [5] that the input state (and so the quantum information it carries) has been nearly perfectly transferred to the bc Hilbert space. What we found therefore is an explicit (and simple) realization of the one-shot state merging protocol [5, 31] (generalizing Ref. [6] studied in the asymptotic limit) and thus the action of $\hat{\mathcal{T}}_{a \rightarrow a'}$ in (1) turns out to be a simple case of the FQSW protocol [7] (the erasure map is essentially the trace map which is a special case of the partial trace appearing in FQSW).

The question of how exactly the information is encoded in the outgoing radiation described by the density matrix ϑ_{bc} is non-trivial. The mapping $\mathcal{T}_{a \rightarrow bc}$ is a highly convoluted identity, something that is expected from the decoupling mechanism. Hence the information is not readily available for a read-out—the state occupying the bc




FIG. 2. Black hole entropy $H_n(t) = H[\vartheta_{a'}] = H[\vartheta_{bc}(t)]$ as a function of the iteration number N for initial states $|n\rangle_a |0\rangle_{bc}$ with $n = 5$ (orange), $n = 20$ (red) and $n = 50$ (pink), with $t = \Delta t N$. Logarithms are base $n + 1$ so that each black hole entropy $H_n(t) \leq 1$. We use $r_0 = 1$ and $\Delta t = 1/15$. The curves are qualitatively unchanged if $\Delta t = 1/25$ and using a commensurately larger number of time slices (data not shown).

Hilbert space is *scrambled*, or, in quantum information theory jargon, it is a highly entangled quantum code.

The output state is highly entangled and so an observer having access only to the b or c mode cannot learn anything. This is a direct consequence of the symmetry between the modes b and c in the boson trilinear Hamiltonian given by Eq. (3), also seen in the action of the unitary $W^{(1)}$ and W in Eqs. (10) and (11), respectively. This is a textbook example of how the impossibility of perfect cloning prevents an observer from obtaining any information from ϑ_b or ϑ_c alone. If there existed a completely positive map that recovers ψ_a from ϑ_b only, the same operation could recover it from ϑ_c . But that would result in two nearly perfect copies of ψ_a which is prohibited by quantum mechanics. Using the language of quantum Shannon theory, the quantum capacity of the channel $\mathcal{M}_{a \rightarrow b(c)}$ must vanish [32].

The trilinear Hamiltonian Eq. (3) that allowed us to carry through this analysis is, to put it bluntly, a guess. It does not follow from an interacting quantum field theory of gravity (such as a locally-flat conformal field theory coupled to a dilaton [33] (see [34] for additional references). However, it is manifestly an elegant extension of the semi-classical approach, and it is difficult to imagine that a consistent renormalizable quantum field theory of gravity would not, in an effective limit, reduce to precisely an interaction as we describe. In fact, the mechanism of information conservation implied by our model is an example of a much larger class of more complicated Hamiltonians with the proper semiclassical limit $V_{\omega_i} |\text{vac}\rangle = 1/\cosh r_{\omega_i} \sum_{m=0}^{\infty} \tanh^m r_{\omega_i} |m\rangle_{b_i} |m\rangle_{c_i}$. Indeed, it may be possible to use decoupling as a constraint to impose conditions on the form of the Bogoliubov transformation that connects the in-vacuum with

the out-vacuum. It is clear from general arguments [35] that at a minimum extra degrees of freedom are needed that are coupled to the Hawking modes, and in our model these degrees of freedom are the quantized black hole modes a .

To illustrate the mechanism concretely, we set $\psi_a = |3\rangle_a$ and observe the behavior of the probabilities given by Eq. (12) in Fig. 3. The black hole state slowly “drifts”




FIG. 3. The schematic evolution of the probability Eq. (12) of the black hole density matrix $\vartheta_{a'}(t)$ for the input state $|3\rangle_a$. Time goes from left, the boxes are Fock bases ($|0\rangle, \dots, |3\rangle$, top to bottom) and the lines represent the contributions from the previous iterations. The vacuum gets the majority of contributions in every iteration (three steps shown here) capturing the black hole evaporation.

towards the vacuum state (the top line) even if the transition amplitudes for the highest Fock state are close to one, that is, the least favorable conditions for the transition to the vacuum state. This is a direct consequence of the spectrum boundedness of the trilinear Hamiltonian Eq. (3) – the number of bosons n in the a mode that $W = \exp[-iH_{\text{tri}}]$ can “distribute” to the bc modes is naturally equal to or lower than n . Crucially, any spectrum-bounded and possibly multilinear Hamiltonian displays exactly the same behavior, and only the information rate differs. As a consequence, the information transfer mechanism—and therefore the Page curve—are expected to remain virtually unchanged even for much more complicated internal dynamics. In other words, the Page curve is robust and universal.

The general features of information transfer between quantum subsystems as described by Page [3] (see also more recently [4, 36, 37]) have been discussed intently in the past. We believe our calculation is the first explicit realization of these previous abstract arguments, pointing towards decoupling as a mechanism for the resolution of the black hole information loss problem. Finally, we would like to point out that the explicitly time-dependent black hole evaporation process we described throws new light on the discussion of putative firewalls [38, 39] for old black holes. The usual line of reasoning for firewalls based on the monogamy property of entanglement does not apply here since the black hole state $|\Psi(t)\rangle_{a'bc}$ from (11) is tripartite entangled.

CA thanks Steve Giddings for discussions. This work was supported in part by Michigan State University through computational resources provided by the Institute for Cyber-Enabled Research.

* kbradler@uottawa.ca

† adami@msu.edu

- [1] A. Fabbri and J. Navarro-Salas, *Modeling Black Hole Evaporation* (Imperial College Press, London (UK), 2005).
- [2] S. W. Hawking, “Breakdown of predictability in gravitational collapse,” *Physical Review D* **14**, 2460 (1976).
- [3] Don N Page, “Information in black hole radiation,” *Physical Review Letters* **71**, 3743 (1993).
- [4] Patrick Hayden and John Preskill, “Black holes as mirrors: quantum information in random subsystems,” *Journal of High Energy Physics* **2007**, 120 (2007).
- [5] Frédéric Dupuis, Mario Berta, Jürg Wullschlegler, and Renato Renner, “One-shot decoupling,” *Communications in Mathematical Physics* **328**, 251–284 (2014).
- [6] Michał Horodecki, Jonathan Oppenheim, and Andreas Winter, “Quantum state merging and negative information,” *Communications in Mathematical Physics* **269**, 107–136 (2007).
- [7] Anura Abeyesinghe, Igor Devetak, Patrick Hayden, and Andreas Winter, “The mother of all protocols: Restructuring quantum information’s family tree,” *Proceedings of the Royal Society of London A: Mathematical, Physical and Engineering Sciences*, 2537–2563 (2009).
- [8] Patrick Hayden, Michał Horodecki, Andreas Winter, and Jon Yard, “A decoupling approach to the quantum capacity,” *Open Systems & Information Dynamics* **15**, 7–19 (2008).
- [9] Frédéric Dupuis, “The decoupling approach to quantum information theory,” arXiv preprint arXiv:1004.1641 (2010).
- [10] U. Leonhardt, *Essential Quantum Optics: From Quantum Measurements to Black Holes* (Cambridge University Press, 2010).
- [11] P. D. Nation, J. R. Johansson, M. P. Blencowe, and Franco Nori, “Colloquium : Stimulating uncertainty: Amplifying the quantum vacuum with superconducting circuits,” *Rev. Mod. Phys.* **84**, 1–24 (2012).
- [12] H. R. Lewis Jr. and W. B. Riesenfeld, “An exact quantum theory of the time-dependent harmonic oscillator and of a charged particle in a time-dependent electromagnetic field,” *J. Math. Phys.* **10**, 1458 (1969).
- [13] I.A. Pedrosa, “Coherent states for certain time-dependent systems,” *Rev. Bras. de Física* **19**, 502–515 (1989).
- [14] G. L. Alberghi, R. Casadio, G. P. Vacca, and G. Venturi, “Gravitational collapse of a radiating shell,” *Phys. Rev. D* **64**, 104012 (2001).
- [15] Tanmay Vachaspati, Dejan Stojkovic, and Lawrence M. Krauss, “Observation of incipient black holes and the information loss problem,” *Phys. Rev. D* **76**, 024005 (2007).
- [16] Anshul Saini and Dejan Stojkovic, “Radiation from a collapsing object is manifestly unitary,” *Phys Rev Lett* **114**, 111301 (2015).
- [17] Stephen W Hawking, “Particle creation by black holes,” *Communications in Mathematical Physics* **43**, 199–220 (1975).
- [18] P. D. Nation and M. P. Blencowe, “The trilinear Hamiltonian: A zero-dimensional model of Hawking radiation from a quantized source,” *New J. Phys.* **12**, 095013 (2010).
- [19] Paul M Alsing, “Parametric down conversion with a depleted pump as a model for classical information transmission capacity of quantum black holes,” *Classical and Quantum Gravity* **32**, 075010 (2015).
- [20] Daniel F Walls and Richard Barakat, “Quantum-mechanical amplification and frequency conversion with a trilinear Hamiltonian,” *Physical Review A* **1**, 446 (1970).
- [21] C Brif, “Coherent states for quantum systems with a trilinear boson Hamiltonian,” *Physical Review A* **54**, 5253 (1996).
- [22] GP Agrawal and CL Mehta, “Dynamics of parametric processes with a trilinear Hamiltonian,” *Journal of Physics A: Mathematical, Nuclear and General* **7**, 607 (1974).
- [23] Markus Aspelmeyer, Tobias J Kippenberg, and Florian Marquardt, “Cavity optomechanics,” *Reviews of Modern Physics* **86**, 1391 (2014).
- [24] U. H. Gerlach, “The mechanism of blackbody radiation from an incipient black hole,” *Phys Rev D* **14**, 1479 (1976).
- [25] P. Arve, G. Bertsch, B. Lauritzen, and G. Puddu, “Static path approximation of the nuclear partition function,” *Annals of Physics* **183**, 309–319 (1988).
- [26] GH Lang, CW Johnson, SE Koonin, and WE Ormand, “Monte Carlo evaluation of path integrals for the nuclear shell model,” *Physical Review C* **48**, 1518–1545 (1993).
- [27] Wilhelm Magnus, “On the exponential solution of differential equations for a linear operator,” *Communications on pure and applied mathematics* **7**, 649–673 (1954).
- [28] Masuo Suzuki, “Generalized Trotter’s formula and systematic approximants of exponential operators and inner derivations with applications to many-body problems,” *Communications in Mathematical Physics* **51**, 183–190 (1976).
- [29] Hale F Trotter, “On the product of semi-groups of operators,” *Proceedings of the American Mathematical Society* **10**, 545–551 (1959).
- [30] Richard P Stanley, *Enumerative combinatorics. Vol. 2* (Cambridge University Press, 1999).
- [31] Mario Berta, “Single-shot quantum state merging,” arXiv preprint arXiv:0912.4495 (2009).
- [32] Kamil Brádler and Christoph Adami, “The capacity of black holes to transmit quantum information,” *Journal of High Energy Physics* **05(2014)095** (2014).
- [33] Jr Callan, Giddings, Harvey, and Strominger, “Evanescent black holes,” *Phys Rev D* **45**, 1005–1009 (1992).
- [34] D. Grumiller, W. Kummer, and D.V. Vassilevich, “Dilaton gravity in two dimensions,” *Phys. Rep.* **369**, 327–340 (2002).
- [35] Sabine Hossenfelder, “Disentangling the black hole vacuum,” *Physical Review D* **91**, 044015 (2015).
- [36] Steven B. Giddings and Yinbo Shi, “Quantum information transfer and models for black hole mechanics,” *Physical Review D* **87**, 064031 (2012).
- [37] Samir D. Mathur and Christopher J. Plumberg, “Correlations in Hawking radiation and the infall problem,” *Journal of High Energy Physics* **JHEP09**, 093 (2011).
- [38] Ahmed Almheiri, Donald Marolf, Joseph Polchinski, and James Sully, “Black holes: complementarity or firewalls?” *Journal of High Energy Physics* **2013**, 1–20 (2013).
- [39] Samuel L Braunstein, Stefano Pirandola, and Karol Życzkowski, “Better late than never: Information re-

trieval from black holes,” *Physical Review Letters* **110**, 101301 (2013).